



US007223552B2

(12) **United States Patent**  
**Hazen et al.**(10) **Patent No.:** **US 7,223,552 B2**(45) **Date of Patent:** **\*May 29, 2007**(54) **MYELOPEROXIDASE, A RISK INDICATOR  
FOR CARDIOVASCULAR DISEASE**(75) Inventors: **Stanley Hazen**, Pepper Pike, OH (US);  
**Renliang Zhang**, Cleveland, OH (US)(73) Assignee: **The Cleveland Clinic Foundation**,  
Cleveland, OH (US)( \* ) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 574 days.This patent is subject to a terminal dis-  
claimer.(21) Appl. No.: **10/039,753**(22) Filed: **Jan. 2, 2002**(65) **Prior Publication Data**

US 2002/0164662 A1 Nov. 7, 2002

**Related U.S. Application Data**(60) Provisional application No. 60/259,340, filed on Jan.  
2, 2001, provisional application No. 60/283,432, filed  
on Apr. 12, 2001.(51) **Int. Cl.****G01N 33/573** (2006.01)**G01N 33/68** (2006.01)(52) **U.S. Cl.** ..... **435/7.24**; 435/7.1; 435/7.4;  
435/28; 436/63; 436/87(58) **Field of Classification Search** ..... 435/7.1,  
435/7.24, 7.4, 28; 436/63, 87  
See application file for complete search history.(56) **References Cited****U.S. PATENT DOCUMENTS**

5,122,534	A	6/1992	Loose et al.
5,731,208	A	3/1998	Heinecke
5,871,946	A *	2/1999	Lucas et al. .... 435/18
5,889,042	A	3/1999	MacLean et al.
5,985,272	A	11/1999	Deby et al.
6,096,556	A	8/2000	Heinecke
6,133,039	A	10/2000	Heinecke
6,268,220	B1	7/2001	Heinecke
6,953,666	B1	10/2005	Kinkade, Jr. et al.
2003/0119792	A1	6/2003	Roca

**FOREIGN PATENT DOCUMENTS**

EP	0 389 381	9/1990
WO	02/48715	6/2002
WO	02/50550	6/2002

**OTHER PUBLICATIONS**"Oxidized LDL and HDL: antagonists in atherothrombosis" by  
Mertens, et al., *FASEB J.*, 15, 2073-2084 (2001)."Macrophage scavenger receptor CD36 is the major receptor for  
LDL modified by monocyte-generated reactive nitrogen species" by  
Podrez, et al., *J. Clin. Invest.* 105:1095-1108 (2000)."Inhibition of Adhesion Molecules Markedly Ameliorates  
Cytokine-Induced Sustained Myocardial Dysfunction in Dogs in  
vivo" by Momii, et al., *J. Mol. Cell. Cardiol.* 30, 2637-2650 (1998)."Supplementation with Tetrahydrobiopterin Suppresses the Devel-  
opment of Hypertension in Spontaneously Hypertension Rats" by  
Hong, et al., *Hypertension*, 2001; 38:1044-1048."Myeloperoxidase Deficiency" by Nauseef, *Hematology, Oncology  
Clinics of North America*, vol. 2, No. 1, Mar. 1988, pp. 135-158."Primary Inherited Defects in Neutrophil Function: Etiology and  
Treatment" by Malech, et al., *Seminars in Hematology*, vol. 34, No.  
4, Oct. 1997, pp. 279-290."Mass spectrometric quantification of amino acid oxidation prod-  
ucts in proteins: insights into pathways that promote LDL oxidation  
in the human artery wall" by Heinecke, et al., *FASEB J.*, 13,  
1113-1120 (1999)."Myeloperoxidase-Generated Oxidants and Atherosclerosis" by  
Podrez, et al., *Free Radical Biology & Medicine*, vol. 28, No. 12, pp.  
1717-1725, Jan. 2000."Myeloperoxidase, a Catalyst for Lipoprotein Oxidation, Is  
Expressed in Human Atherosclerosis Lesions" by Daugherty, et al.,  
*J. Clin. Invest.*, vol. 94, Jul. 1994, 437-444.

(Continued)

*Primary Examiner*—David A. Saunders(74) *Attorney, Agent, or Firm*—Calfee, Halter & Griswold(57) **ABSTRACT**

Diagnostic tests for characterizing an individual's risk of developing or having a cardiovascular disease. In one embodiment the present diagnostic test comprises determining the level of myeloperoxidase (MPO) activity in a bodily sample obtained from the individual or test subject. In another embodiment, the diagnostic test comprises determining the level of MPO mass in a bodily sample obtained from the test subject. In another embodiment, the diagnostic test comprises determining the level of one or more select MPO-generated oxidation products in a bodily sample obtained from the test subject. The select MPO-generated oxidation products are dityrosine, nitrotyrosine, methionine sulphoxide or an MPO-generated lipid peroxidation products. Levels of MPO activity, MPO mass, or the select MPO-generated oxidation product in bodily samples from the test subject are then compared to a predetermined value that is derived from measurements of MPO activity, MPO mass, or the select MPO-generated oxidation product in comparable bodily samples obtained from the general population or a select population of human subjects. Such comparison characterizes the test subject's risk of developing CVD.

**23 Claims, 11 Drawing Sheets**